

Defining the Anthropocene

Simon L. Lewis^{1,2} & Mark A. Maslin¹

Time is divided by geologists according to marked shifts in Earth's state. Recent global environmental changes suggest that Earth may have entered a new human-dominated geological epoch, the Anthropocene. Here we review the historical genesis of the idea and assess anthropogenic signatures in the geological record against the formal requirements for the recognition of a new epoch. The evidence suggests that of the various proposed dates two do appear to conform to the criteria to mark the beginning of the Anthropocene: 1610 and 1964. The formal establishment of an Anthropocene Epoch would mark a fundamental change in the relationship between humans and the Earth system.

Human activity has been a geologically recent, yet profound, influence on the global environment. The magnitude, variety and longevity of human-induced changes, including land surface transformation and changing the composition of the atmosphere, has led to the suggestion that we should refer to the present, not as within the Holocene Epoch (as it is currently formally referred to), but instead as within the Anthropocene Epoch^{1–4} (Fig. 1). Academic and popular usage of the term has rapidly escalated^{5,6} following two influential papers published just over a decade ago^{1,2}. Three scientific journals focusing on the topic have launched: *The Anthropocene*, *The Anthropocene Review* and *Elementa*. The case for a new epoch appears reasonable: what matters when dividing geological-scale time is global-scale changes to Earth's status, driven by causes as varied as meteor strikes, the movement of continents and sustained volcanic eruptions. Human activity is now global and is the dominant cause of most contemporary environmental change. The impacts of human activity will probably be observable in the geological stratigraphic record for millions of years into the future⁷, which suggests that a new epoch has begun⁴.

Nevertheless, some question the types of evidence^{8,9}, because to define a geological time unit, formal criteria must be met^{10,11}. Global-scale changes must be recorded in geological stratigraphic material, such as rock, glacier ice or marine sediments (see Box 1). At present, there is no formal agreement

on when the Anthropocene began, with proposed dates ranging from before the end of the last glaciation to the 1960s. Such different meanings may lead to misunderstandings and confusion across several disciplines. Furthermore, unlike other geological time unit designations, definitions will probably have effects beyond geology. For example, defining an early start date may, in political terms, 'normalize' global environmental change. Meanwhile, agreeing a later start date related to the Industrial Revolution may, for example, be used to assign historical responsibility for carbon dioxide emissions to particular countries or regions during the industrial era. More broadly, the formal definition of the Anthropocene makes scientists arbiters, to an extent, of the human–environment relationship, itself an act with consequences beyond geology. Hence, there is more interest in the Anthropocene than other epoch definitions. Nevertheless, evidence will define whether the geological community formally ratifies a human-activity-induced geological time unit.

We therefore review human geology in four parts. First, we summarize the geologically important human-induced environmental impacts. Second, we review the history of naming the epoch that modern human societies live within, to provide insights into contemporary Anthropocene-related debates. Third, we assess environmental changes caused by human activity that may have left global geological markers consistent with the formal criteria that define geological epochs. Fourth, we highlight the




Figure 1 | Comparison of the current Geologic Time Scale¹⁰ (GTS2012), with two alternatives. a, GTS2012, with boundaries marked in millions of years (ref. 10). b, c, The alternatives include a defined Anthropocene Epoch following either the Holocene (b) or directly following the Pleistocene (c). Defining the Anthropocene as an epoch requires a decision as to whether the Holocene is as distinct as the Anthropocene and Pleistocene; retaining it or not distinguishes between b and c. The question mark represents the current debate over the start of the Anthropocene, assuming it is formally accepted as an epoch (see Box 1, Fig. 2). Colour coding is used according to the Commission for the Geological Map of the World¹⁰, except for the Anthropocene.

¹Department of Geography, University College London, Gower Street, London, WC1E 6BT, UK. ²School of Geography, University of Leeds, Leeds, LS2 9JT, UK.

BOX 1

Dividing geological time

Geological time is divided into a hierarchical series of ever-finer units (Fig. 1a). The present, according to *The Geologic Time Scale 2012*¹⁰, is in the Holocene Epoch (Greek for ‘entirely recent’; started 11,650 yr BP), within the Quaternary Period (started 2.588 million years ago), within the Cenozoic Era (‘recent life’; started 66 million years ago) of the Phanerozoic Eon (‘revealed life’; started 541 million years ago). Divisions represent differences in the functioning of Earth as a system and the concomitant changes in the resident life-forms. Larger differences result in classifications at higher unit-levels.

Formally, geological time units are defined by their lower boundary, that is, their beginning. Boundaries are demarcated using a GSSP, or if good candidate GSSPs do not exist, by an agreed date, termed a GSSA¹⁰. For a GSSP, a ‘stratotype section’ refers to a portion of material that develops over time (rock, sediment, glacier ice), and ‘point’ refers to the location of the marker within the stratotype. Each ‘golden spike’ is a single physical manifestation of a change recorded in a stratigraphic section, often reflecting a global-change phenomenon. GSSP markers are then complemented by a series of correlated changes, also recorded stratigraphically, termed auxiliary stratotypes, indicating widespread changes to the Earth system occurring at that time¹⁰. An exemplary GSSP is the Cretaceous–Paleogene period-level boundary, and the start of the Cenozoic Era, when non-avian dinosaurs declined to extinction and mammals radically increased in variety and abundance. The GSSP boundary marker is the peak in iridium—a residual of bolide impact with Earth—in rock dated at 66 million years ago, located at El Kef, Tunisia¹⁰.

The widespread appearance of new species can also be used as GSSP boundary markers; for example, the Ordovician–Silurian period-level boundary, 443.8 million years ago, is marked by the appearance of a distinct planktonic graptolite, *Akidograptus ascensus* (a now-extinct hemichordate)¹⁰. From an Anthropocene perspective this example shows that the GSSP primary marker chosen as a boundary indicator may be of limited importance compared to the other events taking place that collectively show major changes to Earth at that time⁶⁷.

Formally, a GSSP must have (1) a principal correlation event (the marker), (2) other secondary markers (auxiliary stratotypes), (3) demonstrated regional and global correlation, (4) complete continuous sedimentation with adequate thickness above and below the marker, (5) an exact location—latitude, longitude and height/depth—because a GSSP can be located at only one place on Earth, (6) be accessible, and (7) have provisions for GSSP conservation and protection¹⁰.

Alternatively, following a survey of the stratigraphic evidence, a GSSA date may be agreed by committee to mark a time unit boundary. GSSAs are typical in the Precambrian (>541 million years ago) because well-defined geological markers and clear events are less obvious further back in time¹⁰. Regardless of the marker type, formally ratifying a new Anthropocene Epoch into the GTS would first require a positive recommendation from the Anthropocene Working Group of the Subcommission of Quaternary Stratigraphy, followed by a supermajority vote of the International Commission on Stratigraphy, and finally ratification by the International Union of Geological Sciences¹⁰ (see ref. 11 for full details).

advantages and disadvantages of the few global markers that may indicate a date to define the beginning of the Anthropocene. By consolidating research from disparate fields and the emerging Anthropocene-specific literature we aim to constrain the number of possible Anthropocene start dates, highlight areas requiring further research, and assist in moving towards an evidence-based decision on the possible ratification of a new Anthropocene Epoch.

The geological importance of human actions

Human activity profoundly affects the environment, from Earth’s major biogeochemical cycles to the evolution of life. For example, the early-twentieth-century invention of the Haber–Bosch process, which allows the conversion of atmospheric nitrogen to ammonia for use as fertilizer, has altered the global nitrogen cycle so fundamentally that the nearest suggested geological comparison refers to events about 2.5 billion years ago¹². Human actions have released 555 petagrams of carbon (where 1 Pg = 10¹⁵ g = 1 billion metric tons) to the atmosphere since 1750, increasing atmospheric CO₂ to a level not seen for at least 800,000 years, and possibly several million years^{13,14}, thereby delaying Earth’s next glaciation event¹⁵. The released carbon has increased ocean water acidity at a rate probably not exceeded in the last 300 million years¹⁶.

Human action also affects non-human life. Global net primary productivity appears to be relatively constant¹⁷; however, the appropriation of 25–38% of net primary productivity for human use^{17,18} reduces the amount available for millions of other species on Earth. This land-use conversion to produce food, fuel, fibre and fodder, combined with targeted hunting and harvesting, has resulted in species extinctions some 100 to 1,000 times higher than background rates¹⁹, and probably constitutes the beginning of the sixth mass extinction in Earth’s history¹⁹. Species removals are non-random, with greater losses of large-bodied species from both the land and the oceans. Organisms have been transported around the world, including crops, domesticated animals and pathogens on land. Similarly, boats have transferred organisms among once-disconnected oceans. Such movement has led to a small number of extraordinarily common species, new hybrid species²⁰, and a global homogenization of Earth’s biota. Ostensibly, this change is unique since Pangaea separated about 200 million years ago²¹, but such trans-oceanic exchanges probably have no geological analogue.

Furthermore, human actions may well constitute Earth’s most important evolutionary pressure^{22,23}. The development of diverse products, including antibiotics²², pesticides^{22,24}, and novel genetically engineered organisms²⁴, alongside the movement of species to new habitats²⁵, intense harvesting²³ and the selective pressure of higher air temperatures resulting from greenhouse gas emissions, are all likely to alter evolutionary outcomes^{22–25}. Considered collectively, there is no geological analogue²². Furthermore, given that the lifespan of a species is typically 1–10 million years, the rates of anthropogenic environmental change in the near future may exceed the rates of change encountered by many species in their evolutionary history. Human activity has clearly altered the land surface, oceans and atmosphere, and re-ordered life on Earth.

Historical human geology

Human-related geological time units have a long history²⁶. In 1778 Buffon published an early attempt to describe Earth’s history, allocating a human epoch to be Earth’s seventh and final epoch, paralleling the seven-day creation story²⁷. By the nineteenth century, divine intervention was receding from consideration as a geological force. In 1854 the Welsh geologist and professor of theology, Thomas Jenkyn, appears to have first published the idea of an explicitly evidence-based human geological time unit in a series of widely disseminated geology lessons^{28–30}. He describes the then present day as “the human epoch” based on the likely future fossil record²⁸. In his final lecture he wrote, “All the recent rocks, called in our last lesson Post-Pleistocene, might have been called Anthropozoic, that is, human-life rocks.”²⁹. Similarly, the Reverend Haughton’s 1865 *Manual of Geology* describes the Anthropozoic as the “epoch in which we live”³¹, as did the Italian priest and geologist Antonio Stoppani a decade later³². Meanwhile in the USA, the geology professor James Dwight Dana’s then-popular 1863 *Manual of Geology*³³ extensively refers to the “Age of Mind and Era of Man” as the youngest geological time, as did many of his US contemporaries³⁴.

In 1830 Charles Lyell had proposed that contemporary time be termed the Recent epoch³⁵ on the basis of three considerations: the end of the last glaciation, the then-believed coincident emergence of humans, and the

rise of civilizations^{26,35}. In the 1860s, the French geologist Paul Gervais made Lyell's term international, coining the term Holocene, derived from the Greek for 'entirely recent'. Thus, most nineteenth-century geological textbooks feature humans as part of the definition of the most recent geological time units. Critically, there was little discussion about any of these terms—Recent, Holocene or Anthropozoic—probably because each represented the same conceptual model and broad agreement that humans were part of the definition of the contemporary geological epoch. However, the wider written records of these often deeply religious men show that a separate human epoch was likely to have been more strongly influenced by theological concerns—in particular, separating *Homo sapiens* from other animals and retaining humans at the apex of life on Earth—than by the appraisal of stratigraphic evidence.

In the twentieth century, geologists in the West increasingly used the term Holocene for the current epoch, and Quaternary for the period. Meanwhile, in 1922 the Russian geologist Aleksei Pavlov described the present day as part of an "Anthropogenic system (period) or Anthropocene"³⁶. The Ukrainian geochemist Vladimir Vernadsky then brought to widespread attention the idea that the biosphere, combined with human cognition, had created the Noosphere (from the Greek for mind), with humans becoming a geological force³⁷. The term Noosphere was not well used, but non-Western scientists often used anthropogenic geological time units. The Russian term was anglicized as both Anthropogene and Anthropocene³⁶, sometimes creating confusion. The East–West differences in usage may have been due to differing political ideologies: an orthodox Marxist view of the inevitability of global collective human agency transforming the world politically and economically requires only a modest conceptual leap to collective human agency as a driver of environmental transformation. Again there was little broad interest in the various terms. The Holocene became the official term within the Geologic Time Scale (GTS; Fig. 1)^{10,38}, with its implication that the current interglacial differs from the previous Pleistocene interglacials owing to the influence of humans. It has therefore been argued that an Anthropocene Epoch is not required, given that some human influence is already contained within the definition of the Holocene Epoch⁹. Alternatively, defining the Anthropocene would deprive the Holocene Epoch of its ostensibly unique feature—humans—suggesting that the Holocene as an epoch may not be required.

The views of nineteenth- and twentieth-century scientists illustrate the influence of the dominant contemporary concerns on geological debates. Today's scientists may also not be immune to such influences. For example, a key concern for scientists and others is the central role of technology in modern society and its environmental impacts. Crutzen and Stoermer¹ originally proposed that the start of the Anthropocene should be coincident with the beginning of the Industrial Revolution and James Watt's 1784 refinement of the steam engine. Others followed, including stratigraphers, suggesting that 1800 should be the beginning of the Anthropocene^{39,40}, despite a lack of corresponding global geological markers, and the presence of well-known stratigraphic evidence suggestive of different dates, such as the radionuclide fallout from mid-twentieth-century nuclear weapons tests. Care is needed to ensure that the dominant culture of today's scientists does not subconsciously influence the assessment of stratigraphic evidence.

A human golden spike

Defining the beginning of the Anthropocene as a formal geologic unit of time requires the location of a global marker of an event in stratigraphic material, such as rock, sediment, or glacier ice, known as a Global Stratotype Section and Point (GSSP), plus other auxiliary stratigraphic markers indicating changes to the Earth system. Alternatively, after a survey of the stratigraphic evidence, a date can be agreed by committee, known as a Global Standard Stratigraphic Age (GSSA). GSSPs, known as 'golden spikes', are the preferred boundary markers¹⁰ (see Box 1).

Generally, geologists have used temporally distant changes in multiple stratigraphic records to delimit major changes in the Earth system and thereby geological time units, for example, the appearance of new species as fossils within rocks, coupled with other temporally coincident changes.

Perhaps the most useful GSSP example when considering a possible Anthropocene GSSP is that marking the beginning of the most recent epoch, the Holocene³⁸, because some similar choices and difficulties were faced. These include: not relying on solid aggregate mineral deposits ('rock') for the boundary; an event horizon largely lacking fossils (although fossils are used to recognize Holocene deposits); the need for very precise GSSP dating of events in the recent past; and how to formalize a time unit that extends to the present and thereby implicitly includes a view of the future.

Depending on the parameter considered, the current interglacial took decades to millennia to unfold, as global climate, atmospheric chemistry and the distribution of plant and animal species all altered. From these changes a single dated level within a single stratigraphic record was required to be chosen as a GSSP primary marker (Box 1; Fig. 2). Thus, formally, the Holocene is marked by an abrupt shift in deuterium (²H) excess values at a depth of 1,492.25 m in the NorthGRIP Greenland ice core, dated 11,650 ± 99 yr BP (before present, where 'present' is defined to be 1950)³⁸. This corresponds to the first signs of predominantly Northern Hemisphere climatic warming at the end of the Younger Dryas/ Greenland Stadial 1 cold period³⁸ (Fig. 2). Five further auxiliary stratotypes (four lakes and one marine sediment) showing clear correlated changes across the boundary complement the GSSP, consistent with the occurrence of global changes to the Earth system³⁸. The requirements for a formal definition of the start of the Anthropocene are similar: a clear, datable marker documenting a global change that is recognizable in the stratigraphic record, coupled with auxiliary stratotypes documenting long-term changes to the Earth system.

Defining the Anthropocene presents a further challenge. Changes to the Earth system are not instantaneous. However, even spatially heterogeneous and diachronous (producing similar stratigraphic material varying in age) changes appear near-instantaneous when viewed millions of years after the event, especially as time-lags often fall within the error range of the dating techniques. In contrast, Anthropocene deposits are commonly dated on decadal or annual scales, so that all changes will appear diachronous, to some extent, from today's perspective (but not from far in the future)^{11,41}. Judgement will be required to assess whether the time-lags following events and their significant global impacts are too long to be of use when defining any Anthropocene GSSP.

Several approaches have been put forward to define when the Anthropocene began, including those focusing on the impact of fire⁴², pre-industrial farming^{43–45}, sociometabolism⁴⁶, and industrial technologies^{1,39,40,41,47}, but the relative merits of the evidence for various starting dates have not been systematically assessed against the requirements of a golden spike. Below, we review the major events in human history and pre-history and their impact on stratigraphic records. We focus on continuous stratigraphic material that may yield markers consistent with a GSSP (lake and marine sediments, glacier ice) and on the types of chemical, climatic and biological changes used to denote other epoch boundaries further in the past. We proceed chronologically forward in time, presenting the reason why each event was originally proposed, evaluate the existence of stratigraphic markers, and assess whether the event provides a potential GSSP. The hypotheses and evidence are summarized in Table 1. Following the evidence review we briefly consider the relative merits of the differing events that probably fulfil the GSSP criteria, and assess related GSSA dates.

Pleistocene human impacts

The first major impacts of early humans on their environment was probably the use of fire. Fossil charcoal captures these events from the Early Pleistocene Epoch^{42,48}. However, fires are inherently local events, so they do not provide a global GSSP. The next suggested candidate is the Megafauna Extinction between 50,000 and 10,000 years ago, given that other epoch boundaries have been defined on the basis of extinctions or on the resultant newly emerging species¹⁰. Overall, during the Megafauna Extinction about half of all large-bodied mammals worldwide, equivalent to 4% of all mammal species, were lost⁴⁹. The losses were not evenly distributed: Africa lost 18%, Eurasia lost 36%, North




Figure 2 | Defining the beginning of the Anthropocene. **a**, Current GTS2012 GSSP boundary between the Pleistocene and Holocene³⁸ (dashed line), with global temperature anomalies (relative to the early Holocene average over the period 11,500 BP to 6,500 BP)¹¹² (blue), and atmospheric carbon dioxide composite¹¹³ on the AICC2012 timescale¹¹⁴ (red). **b**, Early Anthropogenic Hypothesis GSSP suggested boundary (dashed line), which posits that early extensive farming impacts caused global environmental changes, defined here by the inflection and lowest level of atmospheric methane (in parts per billion, p.p.b.) from the GRIP ice core⁵⁹ (green), with global temperature anomalies (relative to the average over the period 1961 to 1990)¹¹⁵ (blue), and atmospheric carbon dioxide¹¹³ (red). **c**, Orbis GSSP suggested boundary (dashed line), representing the collision of the Old and New World peoples and homogenization of once distinct biotas, and defined by the pronounced dip in atmospheric carbon dioxide (dashed line) from the Law Dome ice core^{75,76} (blue), with global temperature data anomalies (relative to the average over the period 1961 to 1990)¹¹⁵ (red). **d**, Bomb GSSP suggested boundary (dashed line), characterized by the peak in atmospheric radiocarbon from annual tree-rings (black)¹⁰³ (the $\Delta^{14}\text{C}$ value is the relative difference between the absolute international standard (base year 1950) and sample activity corrected for the time of collection and $\delta^{13}\text{C}$), with atmospheric carbon dioxide from Mauna Loa, Hawaii, post-1958¹¹⁶, and ice core records pre-1958^{75,76} (red), and global temperature anomalies (relative to the average over the period 1961 to 1990)¹¹⁶ (blue).

America lost 72%, South America lost 83%, and Australia lost 88% of their large-bodied mammalian genera^{50,51}. So the Megafauna Extinction was actually a series of events on differing continents at differing times and therefore lacks the required precision for an Anthropocene GSSP marker.

Origins and impacts of farming

The development of agriculture causes long-lasting anthropogenic environmental impacts as it replaces natural vegetation, and thereby increases

species extinction rates, and alters biogeochemical cycles. Agriculture had multiple independent origins: first occurring about 11,000 years ago in southwest Asia, South America and north China; between 6,000–7,000 years ago in Yangtze China and Central America; and 4,000–5,000 years ago in the savanna regions of Africa, India, southeast Asia, and North America⁵². Thus, the increasing presence of fossil pollen from domesticated plants in sediment is too local and lacking in global synchrony to form a GSSP marker. Critically, for the Holocene GSSP, auxiliary markers within stratigraphic material did not include any human-derived markers³⁸, illustrating the lack of anthropogenic impacts at that time. Long-lasting cultural evidence related to agriculture is similarly constrained. Although ceramics are datable and preserved in stratigraphic records (for example, the mineral mullite⁴¹), they appeared in Africa before agriculture, while early southwest Asian farming cultures did not produce ceramics. Similarly, anthropogenically formed soils, derived from intensive farmland management, have also been suggested as a marker of the Anthropocene⁵³. Although these soils are widespread, like vegetation clearance, they are highly diachronous over about 2,000 yr, thus excluding their use as a GSSP marker⁵⁴.

A series of Neolithic revolutions resulted in the majority of *Homo sapiens* becoming agriculturalists to some extent by around 8,000 yr BP, rising to a maximum of about 99% by about 500 yr BP⁴⁶. The Early Anthropogenic Hypothesis posits that the current interglacial was similar to the previous seven interglacial periods until around 8,000 yr BP^{43,55}. By comparison with the closest astronomical analogue of the current interglacial (795,000–780,000 yr BP)⁵⁵, atmospheric CO_2 should have continued to decline after 8,000 yr BP, eventually reaching about 240 parts per million (p.p.m.), and the onset of glaciation should have begun^{43,55}. However, by 6,000–8,000 yr BP, farmers' conversion of high-carbon storage vegetation (forest, woodland, woody savanna) to crops and grazing lands, plus associated fire impacts, may have increased atmospheric CO_2 levels, and postponed this new glaciation⁴³ (Fig. 2). Thus, the lowest level of CO_2 within an ice core record could, in principle, provide a golden spike, but the CO_2 record lacks a distinct inflection point at this time (Fig. 2). Furthermore, the evidence that human activity was responsible for the gradual increase in CO_2 after 6,000 yr BP is extensively debated^{43,56–58}.

Methane provides a clearer inflection point, which may provide a possible GSSP at 5,020 yr BP, the date of the lowest methane value recorded in the GRIP ice core⁵⁹ (Fig. 2). Archaeological evidence suggests that the inflection is caused by rice cultivation in Asia and the expansion of populations of domesticated ruminants. Comparisons of changes in atmospheric methane from the current and past interglacials⁴³, and some methane $\delta^{13}\text{C}$ value evidence⁶⁰, also suggest a human cause. However, a model study suggests that orbital forcing altering methane emissions from tropical wetlands may be responsible⁶¹. Auxiliary markers could include stone axes and fossilized domesticated crop pollen and ruminant remains, but these do not provide temporally well-correlated markers that collectively document globally synchronous changes to the Earth system.

Collision of the Old and New Worlds

The arrival of Europeans in the Caribbean in 1492, and subsequent annexing of the Americas, led to the largest human population replacement in the past 13,000 years⁶², the first global trade networks linking Europe, China, Africa and the Americas^{63,64}, and the resultant mixing of previously separate biotas, known as the Colombian Exchange^{63,64}. One biological result of the exchange was the globalization of human foodstuffs. The New World crops maize/corn, potatoes and the tropical staple manioc/cassava were subsequently grown across Europe, Asia and Africa. Meanwhile, Old World crops such as sugarcane and wheat were planted in the New World. The cross-continental movement of dozens of other food species (such as the common bean, to the New World), domesticated animals (such as the horse, cow, goat and pig, all to the Americas) and human commensals (the black rat, to the Americas), plus accidental transfers (many species of earth worms, to North America; American mink to Europe) contributed to a swift, ongoing, radical reorganization of life on Earth without geological precedent.

Table 1 | Potential start dates for a formal Anthropocene Epoch

Event	Date	Geographical extent	Primary stratigraphic marker	Potential GSSP date*	Potential auxiliary stratotypes
Megafauna extinction	50,000–10,000 yr BP	Near-global	Fossil megafauna	None, diachronous over ~40,000 yr	Charcoal in lacustrine deposits
Origin of farming	~11,000 yr BP	Southwest Asia, becoming global	Fossil pollen or phytoliths	None, diachronous over ~5,000 yr	Fossil crop pollen, phytoliths, charcoal
Extensive farming	~8,000 yr BP to present	Eurasian event, global impact	CO ₂ inflection in glacier ice	None, inflection too diffuse	Fossil crop pollen, phytoliths, charcoal, ceramic minerals
Rice production	6,500 yr BP to present	Southeast Asian event, global impact	CH ₄ inflection in glacier ice	5,020 yr BP CH ₄ minima	Stone axes, fossil domesticated ruminant remains
Anthropogenic soils	~3,000–500 yr BP	Local event, local impact, but widespread	Dark high organic matter soil	None, diachronous, not well preserved	Fossil crop pollen
New–Old World collision	1492–1800	Eurasian–Americas event, global impact	Low point of CO ₂ in glacier ice	1610 CO ₂ minima	Fossil pollen, phytoliths, charcoal, CH ₄ , speleothem $\delta^{18}\text{O}$, tephra†
Industrial Revolution	1760 to present	Northwest Europe event, local impact, becoming global	Fly ash from coal burning	~1900 (ref. 94); diachronous over ~200 yr	¹⁴ N: ¹⁵ N ratio and diatom composition in lake sediments
Nuclear weapon detonation	1945 to present	Local events, global impact	Radionuclides (¹⁴ C) in tree-rings	1964 ¹⁴ C peak§	²⁴⁰ Pu: ²³⁹ Pu ratio, compounds from cement, plastic, lead and other metals
Persistent industrial chemicals	~1950 to present	Local events, global impact	For example, SF ₆ peak in glacier ice	Peaks often very recent so difficult to accurately date§	Compounds from cement, plastic, lead and other metals

For compliance with a Global Stratotype Section and Point (GSSP) definition, a clearly dated global marker is required, backed by correlated auxiliary markers that collectively indicate global and other widespread and long-term changes to the Earth system. BP, before present, where present is defined as calendar date 1950.

* Requires a specific date for a GSSP primary marker. † From Huaynaputina eruption in 1600 (refs 78, 79).

§ Peak, rather than earliest date of detection selected, because earliest dates reflect available detection technology, are more likely influenced by natural background geochemical levels¹⁰¹, and will be more affected by the future decay of the signal, than peak values.

In terms of stratigraphy, the appearance of New World plant species in Old World sediments—and vice versa—may provide a common marker of the Anthropocene across many deposits because pollen is often well preserved in marine and lake sediments. For example, pollen of New World native *Zea mays* (maize/corn), which preserves very well⁴¹, first appears in a European marine sediment core in 1600⁶⁵. The European Pollen Database lists a further 70 lake and marine sediment cores containing *Zea mays* after this date. Phytoliths can similarly record such range expansions⁶⁶. Specifically, the transcontinental range extension of at least one Old World species into the New World (banana, as phytoliths in Central and tropical South America sediments) and a second species from the New World expanding into the Old World (maize/corn, as pollen preserved in sediments in Eurasia and Africa) together constitute a unique signature in the stratigraphic record. This transcontinental range expansion—stratigraphically marking before and after an event—is comparable to the use of the appearance of new species as boundary markers in other epoch transitions^{49,67}.

Besides permanently and dramatically altering the diet of almost all of humanity, the arrival of Europeans in the Americas also led to a large decline in human numbers. Regional population estimates sum to a total of 54 million people in the Americas in 1492⁶⁸, with recent population modelling estimates of 61 million people⁵⁸. Numbers rapidly declined to a minimum of about 6 million people by 1650 via exposure to diseases carried by Europeans, plus war, enslavement and famine^{58,63,68,69}. The accompanying near-cessation of farming and reduction in fire use resulted in the regeneration of over 50 million hectares of forest, woody savanna and grassland with a carbon uptake by vegetation and soils estimated at 5–40 Pg within around 100 years^{58,70–72}. The approximate magnitude and timing of carbon sequestration suggest that this event significantly contributed to the observed decline in atmospheric CO₂ of 7–10 p.p.m. (1 p.p.m. CO₂ = 2.1 Pg of carbon) between 1570 and 1620 documented in two high-resolution Antarctic ice core records^{73–76} (Fig. 2 and Box 2). This dip in atmospheric CO₂ is the most prominent feature, in terms of both rate of change and magnitude, in pre-industrial atmospheric CO₂ records over the past 2,000 years⁷⁵ (Fig. 2).

On the basis of the movement of species, atmospheric CO₂ decline and the resulting climate-related changes within various stratigraphic records, we propose that the 7–10 p.p.m. dip in atmospheric CO₂ to a

low point of 271.8 p.p.m. at 285.2 m depth of the Law Dome ice core⁷⁵, dated 1610 (± 15 yr; refs 75, 76), is an appropriate GSSP marker (Fig. 2). Auxiliary stratotypes could include: the first occurrence of a cross-ocean range extension in the fossil record (*Zea mays*, in 1600⁶⁵) plus a range of deposits showing distinct changes at that time, including tephra^{77,78} and other signatures from the 1600 Huaynaputina eruption detected at both poles and in the tropics^{77–79}; charcoal reductions in deposits in the Americas⁷¹ and globally⁸⁰; decreases in atmospheric methane, enrichment of methane $\delta^{13}\text{C}$, and decreases in carbon monoxide in Antarctic ice cores^{60,81–84}; pollen in lacustrine sediments showing vegetation regeneration⁸⁵; proxies indicating anomalous Arctic sea-ice extent⁸⁶; changing $\delta^{18}\text{O}$ derived from speleothems from caves in China and Peru¹⁴ and other studies noting changes coincident with 1600 and the coolest part of the Little Ice Age (1594–1677; ref. 87), a relatively synchronous global event noted in geologic deposits worldwide⁸⁷.

The impacts of the meeting of Old and New World human populations—including the geologically unprecedented homogenization of Earth's biota^{63,64}—may serve to mark the beginning of the Anthropocene. Although it represents a major event in world history^{62–64,88}, the collision of the Old and New Worlds has not been proposed previously, to our knowledge, as a possible GSSP. We suggest naming the dip in atmospheric CO₂ the 'Orbis spike' and the suite of changes marking 1610 as the beginning of the Anthropocene the 'Orbis hypothesis', from the Latin for world, because post-1492 humans on the two hemispheres were connected, trade became global, and some prominent social scientists refer to this time as the beginning of the modern 'world-system'⁸⁹.

Industrialization

The beginning of the Industrial Revolution has often been suggested as the beginning of the Anthropocene, because accelerating fossil fuel use and coupled rapid societal changes herald something important and unique in human history^{1–4,39}. Yet humans have long been engaging in industrial-type production, such as metal utilization from around 8,000 yr BP onwards, with attendant pollution⁹⁰. Elevated mercury records are documented at around 3,400 yr BP in the Peruvian Andes⁹¹, while the impacts of Roman Empire copper smelting are detectable in a Greenland ice core at around 2,000 yr BP⁹². This metal pollution, like other examples predating the Industrial Revolution, is too local and diachronous to provide a golden spike.

BOX 2

Origins of the 1610 decrease in atmospheric CO₂Is the CO₂ decline real?

Two independent high-resolution Antarctic ice core records from the Law Dome and the Western Antarctic Ice Sheet show a reduction in atmospheric CO₂ of 7–10 p.p.m. between 1570 and 1620^{73–75} (Fig. 2). A smaller CO₂ decrease is also observed in less highly resolved Antarctic cores^{117,118}. The decline exceeds the measurement error of the cores, 1–2 p.p.m., and experiments suggest that it does not result from *in situ* changes within the ice core¹¹⁹.

Did human activity cause the decline?

The arrival of Europeans in the Americas led to a catastrophic decline in human numbers, with about 50 million deaths between 1492 and 1650, according to several independent sources^{58,63,68,69}. Contemporary field observations of soil¹²⁰ and vegetation¹²¹ carbon dynamics following agriculture abandonment suggest that about 65 million hectares (that is, 50 million people × 1.3 hectares per person) would sequester 7–14 Pg of carbon over 100 years (that is, 100–200 Mg of carbon per hectare total uptake, above- and below-ground). Reduction in fire use for land management would additionally increase carbon uptake outside farmed areas. Studies using a variety of methods report broadly consistent estimates^{58,70–72} of carbon uptake by vegetation of 5–40 Pg (2.1 Pg of carbon = 1 p.p.m. atmospheric CO₂ over shorter timescales, lessening over time¹²⁷). Given that maximum human mortality rates were not reached for some decades after 1492^{62,63}, and maximum carbon uptake would take place 20–50 yr after farming abandonment, peak carbon sequestration would occur approximately between 1550 and 1650.

Some model studies spanning thousands of years find a net land surface carbon uptake spanning 1500–1650 across the Americas⁵⁸, while others do not¹²². However, in general, evidence from such studies weakly constrain the problem because Holocene carbon cycle modelling is designed to investigate changes associated with long-acting slow processes (carbon uptake by peat or coral reefs) and feedback mechanisms (oceanic outgassing, oceanic uptake and CO₂ fertilization of vegetation), and probably poorly represent the short period of the CO₂ dip (for example, ref. 57). For example, a study calculating a net zero impact of the cessation of farming in the Americas¹²² included a large soil carbon flux to the atmosphere, which contradicts field evidence^{120,123}, and had the effect of offsetting the uptake from growing trees¹²². Carbon cycle models with robust representations of land-use change and subsequent vegetation regeneration following the Americas population catastrophe will be required to improve estimates of carbon uptake compared with carbon accounting studies.

The approximate magnitude and timing of carbon sequestration make the population decline in the Americas the most likely cause of the observed decline in atmospheric CO₂. Atmospheric^{74,124,125} and tropical marine $\delta^{13}\text{C}$ analyses¹²⁶ also support uptake of CO₂ by vegetation rather than oceanic uptake. The 1600 Huaynaputina eruption in Peru^{78,79} probably exacerbated the CO₂ minima, and a lagged oceanic outgassing in response to the land carbon uptake probably contributed to the fast rebound of atmospheric CO₂ after 1610¹²⁷. In addition, multi-proxy reconstructions of temperature indicate that, after accounting for both solar and volcanic radiative forcing, additional terrestrial carbon uptake is required to explain temperature declines over the 1550–1650 period¹⁰⁷. This is consistent with uptake by vegetation following the population crash in the Americas¹⁰⁷.

Definitions of the Industrial Revolution give an onset date anywhere between 1760 and 1880, beginning as an event local to northwest Europe⁸⁸. Given the initial slow spread of coal use, ice core records show little impact on global atmospheric CO₂ concentration until the nineteenth century, and then they show a relatively smooth increase rather than an abrupt change, precluding this as a GSSP marker (Fig. 2). Similarly, other associated changes, including methane and nitrate¹⁵, products of fossil fuel burning (including spherical carbonaceous particles⁹³ and magnetic fly ash⁹⁴) plus resultant changes in lake sediments^{95,96} alter slowly as the use of fossil fuels increased over many decades. Lead, which was once routinely added to vehicle fuels, has been proposed as a possible marker, because leaded fuel was almost globally used and is now banned⁹⁷. However, peak lead isotope ratio values from this source in sediments and other deposits vary from 1940 to after 1980, limiting the utility of this marker. The Industrial Revolution thus provides a number of markers spreading from northwest Europe to North America and expanding worldwide since about 1800, although none provides a clear global GSSP primary marker.

The Great Acceleration

Since the 1950s the influence of human activity on the Earth system has increased markedly. This 'Great Acceleration' is marked by a major expansion in human population, large changes in natural processes^{3,12,98}, and the development of novel materials from minerals to plastics to persistent organic pollutants and inorganic compounds^{41,47,97}. Among these many changes the global fallout from nuclear bomb tests has been proposed as a global event horizon marker^{41,47}. The first detonation was in 1945, with a peak in atmospheric testing from the late 1950s to early 1960s, followed by a rapid decline following the Partial Test Ban Treaty in 1963 and later agreements, such that only low test levels continue to the present day (Fig. 2). A resulting distinct peak in radioactivity is recorded in high-resolution ice cores, lake and salt marsh sediments, corals, speleothems and tree-rings from the early 1950s onwards, declining in the late 1960s^{15,99}. The clearest signal is from atmospheric ¹⁴C, seen in direct air measurements and captured by tree-rings and glacier ice, which reaches a maximum in the mid- to high-latitude Northern Hemisphere at 1963–64 and a year later in the tropics¹⁰⁰. Although ¹⁴C has a relatively short half-life (5,730 years), elevated levels will persist long enough to be useable for several generations of geologists in the future.

While recognizing that many apparently novel industrially produced chemicals are occasionally produced in small quantities naturally¹⁰¹, chemical signatures from long-lived well-mixed gases in glacier ice or sediments may also meet GSSP criteria. Potential long-lived gases are the halogenated gases, such as SF₆, C₂F₆, CF₄ (with half-lives of 3,000 yr, 10,000 yr and 50,000 yr, respectively). Most were first manufactured industrially in the 1950s, and many are measurable in firn air¹⁰², and with large enough samples could be measured in ice cores¹⁵. But although they are measurable, distinct peaks are very recent and sometimes absent because major declines in industrial production are occurring after the negotiation and ratification of the 1989 Montreal and 2005 Kyoto protocols.

Of the various possible mid- to late-twentieth-century markers of the Great Acceleration, the global ¹⁴C peak provides an unambiguously global change in a number of stratigraphic deposits. We suggest that an unequivocally annual record is the optimal choice to reflect the ¹⁴C peak, thereby giving a dating accuracy of one year. We propose that the GSSP marker should be the ¹⁴C peak, at 1964, within dated annual rings of a pine tree (*Pinus sylvestris*) from King Castle, Niepołomice, 25 km east of Kraków, Poland¹⁰³ (Fig. 2). Secondary correlated markers would include plutonium isotope ratios (²⁴⁰Pu/²³⁹Pu) in sediments indicating bomb testing¹⁰⁴, (fast-decaying) 137-Caesium⁹⁷, alongside the presence of peaks in very long-lived iodine isotopes (¹²⁹I, with half-life 15.7 million years) found in marine sediments¹⁰⁵ and soils¹⁰⁶.

While radionuclide fallout did not have major biological or other widespread physical repercussions, other auxiliary stratotypes may include the numerous other human-driven changes resulting in mid- to late-twentieth-century changes in geological deposits, including fossil pollen of novel genetically modified crops; declines in $\delta^{15}\text{N}$ in Northern Hemisphere

lakes⁹⁶ and ice cores¹⁵; the emergence of SF₆ and CF₄ from background levels¹⁵; lead isotopes in ice cores¹⁵; microplastics in marine sediments⁹⁷; diatom assemblages in lakes in response to eutrophication⁴¹; and benthic foraminifera changes in marine sediments⁴¹.

Dating the Anthropocene

We conclude that most proposed Anthropocene start dates, including the earliest detectable human impacts⁴², earliest widespread impacts⁴⁵, and historic events such as the Industrial Revolution^{1–3,39,40}, can probably be rejected because they are not derived from a globally synchronous marker. Our review highlights that only those environmental changes associated with well-mixed atmospheric gases provide clearly global synchronous geological markers on an annual or decadal scale, as is required to define a GSSP for the Anthropocene. The earliest potential GSSP primary marker we identify is the inflection of atmospheric methane at 5,020 yr BP (Fig. 2; Table 1), but correlated auxiliary stratotypes are lacking. Thus, the CH₄ inflection is unlikely to be a strong candidate for the beginning of the Anthropocene. We find that only two other events—the Orbis spike dip in CO₂ with a minimum at 1610, and the bomb spike 1964 peak in ¹⁴C—appear to fulfil the criteria for a GSSP to define the inception of the Anthropocene (Fig. 2; Table 1). While both GSSP dates have a number of correlated auxiliary stratotypes there are advantages and disadvantages associated with each.

The main advantage to the 1610 Orbis spike is the geological and historical importance of the event. In common with other epoch boundaries¹⁰ this boundary would document changes in climate^{87,107}, chemistry⁷⁵ and palaeontological^{65,85} signals. Critically, the transoceanic movement of species is an unambiguously permanent change to the Earth system⁴⁰, and such a boundary would mark Earth's last globally synchronous cool period⁸⁷ before the long-term global warmth of the Anthropocene Epoch. Historically, the Industrial Revolution has often been considered as the most important event in relation to the inception of the Anthropocene^{1,2,39,40}, but we have not identified a clear global Industrial Revolution GSSP. However, in the view of many historians, industrialization and extensive fossil fuel use were only made possible by the annexing of the Americas⁸⁸. Before the Industrial Revolution both northwest Europe and southern China were similar in terms of life expectancy and material consumption patterns, including modest coal use, and both regions faced productive boundaries based on the available land area⁸⁸. Thus, the agricultural commodities from the vast new lands of the Americas allowed Europe to transcend its ecological limits and sustain economic growth. In turn, this freed labour, allowing Europe to industrialize. That is, the Americas made industrialization possible owing to the unprecedented inflow of new cheap resources (and profitable new markets for manufactured goods). This 'Great Divergence' of Europe from the rest of the world required access to and exploitation of new lands plus a rich source of easily exploitable energy: coal⁸⁸. Thus, dating the Anthropocene to start about 150 years before the beginning of the Industrial Revolution is consistent with a contemporary understanding of the likely material causes of the Industrial Revolution. The main disadvantage to the Orbis hypothesis is that a number of deposits may not show large changes around 1600, particularly in terms of biological material from the transport of species to new continents or oceans, because there are time-lags before species newly appear in geological deposits.

The key advantage of selecting 1964 as the base of a new Anthropocene Epoch is the sheer variety of human impacts recorded during the Great Acceleration: almost all stratigraphic records today, and over recent decades, have some marker of human activity. The latter part of the twentieth century is unambiguously a time of major anthropogenic global environmental impacts¹⁰⁸. One disadvantage is that although nuclear explosions have the capacity to fundamentally transform many aspects of Earth's functioning, so far they have not done so, making the radionuclide spike a good GSSP marker but not an Earth-changing event. A further possible limitation in selecting such a recent date is that some deposits, notably some marine sediments, do not accumulate and stabilize over time spans

as short as the past 50 years, making clear datable changes and correlation among some stratotypes sometimes difficult to discern⁴⁰.

Choosing between the 1610 Orbis and 1964 bomb spikes is challenging. As an alternative, a GSSA date, based on stratigraphic evidence, could be agreed upon by committee as the inception of the Anthropocene. However, any chosen date would be potentially open to challenge as arbitrary. For example, the Industrial Revolution is certainly a pivotal moment in human history, yet it is unclear how one could choose, based on the available geological evidence, an early Industrial Revolution GSSA date, say 1800, over a later date, perhaps 1850 or 1900. Similarly, the Great Acceleration is diachronous¹⁰⁸, and GSSA suggested dates could be 1945, 1950 or 1954 (ref. 109). Given such difficulties, given that GSSP markers are preferred¹⁰, and given that candidate GSSP markers exist, a GSSA date seems unnecessary. Of the GSSP possibilities we tend to prefer 1610, because the transoceanic movement of species is a clear and permanent geological change to the Earth system. This date also fits more closely with Crutzen and Stoermer's original proposal¹ of an important historical juncture—the Industrial Revolution—as the beginning of the Anthropocene, which has been enduringly popular and useful, suggesting 1610 may be similarly so.

We hope that identifying a limited number of possible events and GSSP markers may assist in focusing research efforts to select a robust GSSP alongside a series of auxiliary stratotypes. Such research might include compiling data sets of the first appearance of non-native species in lake and marine sediments to better document the transoceanic spread of species and improve the evidence on which the 1610 proposal is based. The reliable detection of ¹²⁹I in high-resolution glacier ice and expanding the number of locations at which novel minerals, compounds and other recent human signals are investigated^{41,47} would advance the 1964 GSSP proposal.

Ratification of an Anthropocene Epoch would require a further decision to be made, that is, whether to retain the Holocene Epoch (Fig. 1). All Anthropocene GSSP choices would leave a complete Holocene Epoch at least three orders of magnitude shorter than any other epoch¹⁰ and similar to previous Pleistocene interglacials⁵⁵, which are not epoch-level events. Furthermore, the existence of a Holocene Epoch is due, in part, to the view—originating from nineteenth-century geologists—that the presence or influence of humans distinguished the Holocene from the Pleistocene^{9,26,27,35,38}. An Anthropocene Epoch, combined with today's evidence that *Homo sapiens* is a Pleistocene species, removes key justifications for retaining the Holocene as an epoch-level designation. We therefore suggest that if the Anthropocene is accepted as an epoch it should directly follow the Pleistocene (Fig. 1c), as suggested independently elsewhere¹¹⁰. If the Holocene ceases to be an epoch but refers instead to the final stage of the Pleistocene Epoch, we suggest that the term Holocenian Stage is used, to maintain consistency with current terminology. While an alternative informal geological term, the Flandrian stage, denotes the current interglacial as part of the Pleistocene, its use has strongly declined over recent decades¹⁰, and would not be as recognizable as the Holocenian Stage. Re-classifying any pre-Anthropocene Epoch interglacial time unit as the Holocenian Stage will create the usual tension¹⁰ between resistance to altering past GTS agreements and the maintenance of GTS internal consistency.

The wider importance

The choice of either 1610 or 1964 as the beginning of the Anthropocene would probably affect the perception of human actions on the environment. The Orbis spike implies that colonialism, global trade and coal brought about the Anthropocene. Broadly, this highlights social concerns, particularly the unequal power relationships between different groups of people, economic growth, the impacts of globalized trade, and our current reliance on fossil fuels. The onward effects of the arrival of Europeans in the Americas also highlights a long-term and large-scale example of human actions unleashing processes that are difficult to predict or manage. Choosing the bomb spike tells a story of an elite-driven technological development that threatens planet-wide destruction. The

long-term advancement of technology deployed to kill people, from spears to nuclear weapons, highlights the more general problem of 'progress traps'¹¹¹. Conversely, the 1963 Partial Test Ban Treaty and later agreements highlight the ability of people to collectively successfully manage a major global threat to humans and the environment. The event or date chosen as the inception of the Anthropocene will affect the stories people construct about the ongoing development of human societies.

Past scientific discoveries have tended to shift perceptions away from a view of humanity as occupying the centre of the Universe. In 1543 Copernicus's observation of the Earth revolving around the Sun demonstrated that this is not the case. The implications of Darwin's 1859 discoveries then established that *Homo sapiens* is simply part of the tree of life with no special origin. Adopting the Anthropocene may reverse this trend by asserting that humans are not passive observers of Earth's functioning. To a large extent the future of the only place where life is known to exist is being determined by the actions of humans. Yet, the power that humans wield is unlike any other force of nature, because it is reflexive and therefore can be used, withdrawn or modified. More widespread recognition that human actions are driving far-reaching changes to the life-supporting infrastructure of Earth may well have increasing philosophical, social, economic and political implications over the coming decades.

Received 26 March 2014; accepted 12 January 2015.

1. Crutzen, P. J. & Stoermer, E. F. The Anthropocene. *IGBP Global Change Newsl.* **41**, 17–18 (2000).
This paper suggested that the Holocene has ended and the Anthropocene has begun, starting the contemporary increase in the usage of the term Anthropocene.
2. Crutzen, P. J. Geology of mankind. *Nature* **415**, 23 (2002).
3. Steffen, W., Crutzen, P. J. & McNeill, J. R. The Anthropocene: are humans now overwhelming the great forces of nature. *Ambio* **36**, 614–621 (2007).
4. Zalasiewicz, J., Williams, M., Haywood, A. & Ellis, M. The Anthropocene: a new epoch of geological time? *Phil. Trans. R. Soc. Lond. A* **369**, 835–841 (2011).
5. Dalby, S. Biopolitics and climate security in the Anthropocene. *Geoforum* **49**, 184–192 (2013).
6. Anon. The Anthropocene: a man-made world. *The Economist* **May 26** (2011); <http://www.economist.com/node/18741749>.
7. Zalasiewicz, J. *The Earth After Us: What Legacy Will Humans Leave in the Rocks?* (Oxford University Press, 2008).
8. Autin, W. J. & Holbrook, J. M. Is the Anthropocene an issue of stratigraphy or pop culture? *GSA Today* **22**, 60–61 (2012).
9. Gibbard, P. L. & Walker, M. J. C. The term 'Anthropocene' in the context of formal geological classification. *Geol. Soc. Lond. Spec. Publ.* **395**, 29–37 (2014).
This paper presents a view that there is not currently enough evidence to formally ratify a new Anthropocene Epoch.
10. Gradstein, F. M., Ogg, J. G., Schmitz, M. D. & Ogg, G. M. *The Geologic Time Scale 2012* (Elsevier, 2012).
This book is the latest GTS, including the formal assessments of Earth's history divided into epochs, periods, eras and eons.
11. Finney, S. C. The 'Anthropocene' as a ratified unit in the ICS International Chronostratigraphic Chart: fundamental issues that must be addressed by the Task Group. *Geol. Soc. Lond. Spec. Publ.* **395**, 23–28 (2014).
This paper details the requirements and questions that will need to be addressed by the initial committee that will recommend whether or not an Anthropocene epoch is to be formally defined.
12. Canfield, D. E., Glazer, A. N. & Falkowski, P. G. The evolution and future of Earth's nitrogen cycle. *Science* **330**, 192–196 (2010).
13. Ciais, P. et al. in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker, T. F. et al.) Ch. 6, 465–570 (Cambridge Univ. Press, 2013).
14. Masson-Delmotte, V. et al. in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker, T. F. et al.) Ch. 5, 383–464 (Cambridge Univ. Press, 2013).
15. Wolff, E. W. Ice Sheets and the Anthropocene. *Geol. Soc. Lond. Spec. Publ.* **395**, 255–263 (2014).
16. International Geosphere-Biosphere Programme, Intergovernmental Oceanographic Commission, Scientific Committee on Oceanic Research. *Ocean Acidification Summary for Policymakers – Third Symposium on the Ocean in a High-CO₂ World* (International Geosphere-Biosphere Programme, 2013), <http://ocean-acidification.net/for-policymakers/>.
17. Running, S. W. A measurable planetary boundary for the biosphere. *Science* **337**, 1458–1459 (2012).
18. Krausmann, F. et al. Global human appropriation of net primary production doubled in the 20th century. *Proc. Natl Acad. Sci. USA* **110**, 10324–10329 (2013).
19. Barnosky, A. D. et al. Has the Earth's sixth mass extinction already arrived? *Nature* **471**, 51–57 (2011).
20. Thomas, C. D. The Anthropocene could raise biological diversity. *Nature* **502**, 7 (2013).
21. Baiser, B., Olden, J. D., Record, S., Lockwood, J. L. & McKinney, M. L. Pattern and process of biotic homogenization in the New Pangaea. *Proc. R. Soc. Lond. B* **279**, 4772–4777 (2012).
22. Palumbi, S. R. Humans as the world's greatest evolutionary force. *Science* **293**, 1786–1790 (2001).
23. Darimont, C. T. et al. Human predators outpace other agents of trait change in the wild. *Proc. Natl Acad. Sci. USA* **106**, 952–954 (2009).
24. Tabashnik, B. E., Mota-Sanchez, D., Whalon, M. E., Hollingworth, R. M. & Carriere, Y. Defining terms for proactive management of resistance to Bt crops and pesticides. *J. Econ. Entomol.* **107**, 496–507 (2014).
25. Stuart, Y. E. et al. Rapid evolution of a native species following invasion by a congener. *Science* **346**, 463–466 (2014).
26. Davis, R. V. Inventing the present: historical roots of the Anthropocene. *Earth Sci. Hist.* **30**, 63–84 (2011).
This paper investigates and reviews the history of the use of the terms 'Holocene' and 'Anthropocene', showing that the Holocene includes humans in its first nineteenth-century definition.
27. Rudwick, M. S. J. *Bursting the Limits of Time: The Reconstruction of Geohistory in the Age of Revolution* (University of Chicago Press, 2005).
28. Jenkyn, T. W. Lessons in Geology XLVI. Chapter IV. On the effects of organic agents on the Earth's crust. *Popular Educator* **4**, 139–141 (1854).
29. Jenkyn, T. W. Lessons in Geology XLIX. Chapter V. On the classification of rocks section IV. On the tertiary. *Popular Educator* **4**, 312–316 (1854).
30. Hansen, P. H. *The Summits of Modern Man: Mountaineering after the Enlightenment* (Harvard University Press, 2013).
31. Houghton, S. *Manual of Geology* (Longman, 1865).
32. Stoppani, A. *Corso di Geologia* Vol. II (G. Bernardoni e G. Brigola, 1873).
33. Dana, J. D. *Manual of Geology* (Theodore Bliss and Co., 1863).
34. Le Conte, J. On critical periods in the history of the Earth and their relation to evolution; and on the Quaternary as such a period. *Am. J. Sci.* **14**, 99–114 (1877).
35. Lyell, C. *Principles of Geology* Volumes I, II and III (University of Chicago Press, 1990); originally published by John Murray, 1830–1833.
36. Shantser, E. V. in *Great Soviet Encyclopedia* Vol. 2 (ed. Prokhorov, A. M.) 139–144 (Macmillan, 1979).
37. Vernadsky, W. I. Biosphere and Noosphere. *Am. Sci.* **33**, 1–12 (1945).
38. Walker, M. et al. Formal definition and dating of the GSSP (Global Stratotype Section and Point) for the base of the Holocene using the Greenland NGRIP ice core, and selected auxiliary records. *J. Quat. Sci.* **24**, 3–17 (2009).
39. Steffen, W., Grinevald, J., Crutzen, P. & McNeill, J. The Anthropocene: conceptual and historical perspectives. *Phil. Trans. R. Soc. Lond. A* **369**, 842–867 (2011).
40. Zalasiewicz, J. et al. Stratigraphy of the Anthropocene. *Phil. Trans. R. Soc. Lond. A* **369**, 1036–1055 (2011).
41. Waters, C. N., Zalasiewicz, J. A., Williams, M., Ellis, M. A. & Snelling, A. M. A stratigraphical basis for the Anthropocene? *Geol. Soc. Lond. Spec. Publ.* **395**, 1–21 (2014).
This paper reviews various stratigraphic markers relevant to defining the Anthropocene, with an up-to-date collation of the many markers coincident with the Industrial Revolution and the Great Acceleration.
42. Glikson, A. Fire and human evolution: the deep-time blueprints of the Anthropocene. *Anthropocene* **3**, 89–92 (2013).
43. Ruddiman, W. F. The Anthropocene. *Annu. Rev. Earth Planet. Sci.* **41**, 45–68 (2013).
This paper summarizes the data and arguments that human activity altered CO₂ and CH₄ emissions thousands of years ago, leading to a delayed next glaciation, known as the Early Anthropogenic Hypothesis.
44. Foley, S. F. et al. The Palaeoanthropocene—the beginnings of anthropogenic environmental change. *Anthropocene* **3**, 83–88 (2013).
45. Balter, M. Archaeologists say the 'Anthropocene' is here—but it began long ago. *Science* **340**, 261–262 (2013).
46. Fischer-Kowalski, M., Krausmann, F. & Pallua, I. A sociometabolic reading of the Anthropocene: modes of subsistence, population size and human impact on Earth. *Anthropocene Rev.* **1**, 8–33 (2014).
This paper takes an alternative view of the Anthropocene, considering human energy sources, and posits two transitions, to an agricultural mode, about 10,000 yr BP, and to an industrial mode, which begins after 1500.
47. Zalasiewicz, J., Williams, M. & Waters, C. N. Can an Anthropocene series be defined and recognized? *Geol. Soc. Lond. Spec. Publ.* **395**, 39–53 (2014).
48. Roebroeks, W. & Villa, P. On the earliest evidence for habitual use of fire in Europe. *Proc. Natl Acad. Sci. USA* **108**, 5209–5214 (2011).
49. Barnosky, A. D. Palaeontological evidence for defining the Anthropocene. *Geol. Soc. Lond. Spec. Publ.* **395**, 149–165 (2014).
50. Barnosky, A. D., Koch, P. L., Feranec, R. S., Wing, S. L. & Shabel, A. B. Assessing the causes of Late Pleistocene extinctions on the continents. *Science* **306**, 70–75 (2004).
51. Lorentzen, E. D. et al. Species-specific responses of Late Quaternary megafauna to climate and humans. *Nature* **479**, 359–364 (2011).
52. Ellis, E. C. et al. Used planet: a global history. *Proc. Natl Acad. Sci. USA* **110**, 7978–7985 (2013).
53. Certini, G. & Scalenghe, R. Anthropogenic soils are the golden spikes for the Anthropocene. *Holocene* **21**, 1269–1274 (2011).
54. Gale, S. J. & Hoare, P. G. The stratigraphic status of the Anthropocene. *Holocene* **22**, 1491–1494 (2012).

55. Tzedakis, P. C., Channell, J. E. T., Hodell, D. A., Kleiven, H. F. & Skinner, L. C. Determining the natural length of the current interglacial. *Nature Geosci.* **5**, 138–141 (2012).
56. Broecker, W. C. & Stocker, T. F. The Holocene CO₂ rise: Anthropogenic or natural? *Eos* **87**, 27–29 (2006).
57. Stocker, B. D., Strassmann, K. & Joos, F. Sensitivity of Holocene atmospheric CO₂ and the modern carbon budget to early human land use: analyses with a process-based model. *Biogeosciences* **8**, 69–88 (2011).
58. Kaplan, J. O. *et al.* Holocene carbon emissions as a result of anthropogenic land cover change. *Holocene* **21**, 775–791 (2011).
59. Blunier, T., Chappellaz, J., Schwander, J., Stauffer, B. & Raynaud, D. Variations in atmospheric methane concentration during the Holocene epoch. *Nature* **374**, 46–49 (1995).
60. Sapart, C. J. *et al.* Natural and anthropogenic variations in methane sources during the past two millennia. *Nature* **490**, 85–88 (2012).
61. Singarayer, J. S., Valdes, P. J., Friedlingstein, P., Nelson, S. & Beerling, D. J. Late Holocene methane rise caused by orbitally controlled increase in tropical sources. *Nature* **470**, 82–85 (2011).
62. Diamond, J. *Guns, Germs and Steel: A Short History of Everybody for the Last 13,000 Years* (Chatto and Windus, 1997).
63. Mann, C. C. 1493: *How the Ecological Collision of Europe and the Americas Gave Rise to the Modern World* (Granta, 2011).
64. Crosby, A. W. *The Columbian Exchange: Biological and Cultural Consequences of 1492* 30 yr edn (Preager, 2003).
65. Mercuri, A. M. *et al.* A marine/terrestrial integration for mid-late Holocene vegetation history and the development of the cultural landscape in the Po valley as a result of human impact and climate change. *Vegetat. Hist. Archaeobot.* **21**, 353–372 (2012).
66. Piperno, D. R. Identifying crop plants with phytoliths (and starch grains) in Central and South America: a review and an update of the evidence. *Quat. Int.* **193**, 146–159 (2009).
67. Zalasiewicz, J. & Williams, M. The Anthropocene: a comparison with the Ordovician-Silurian boundary. *Rendiconti Lincei-Scienze Fisiche E Naturali* **25**, 5–12 (2014).
68. Denevan, W. M. *The Native Population of the Americas in 1492* 2nd edn (University of Wisconsin Press, 1992).
69. Mann, C. C. 1491: *New Revelations of the Americas Before Columbus* (Vintage, 2005).
70. Nevle, R. J. & Bird, D. K. Effects of syn-pandemic fire reduction and reforestation in the tropical Americas on atmospheric CO₂ during European conquest. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **264**, 25–38 (2008).
- This paper presents a synthesis of data computing the impacts of the rapid 1492–1650 reduction in population across the Americas and the carbon uptake implications.**
71. Dull, R. A. *et al.* The Columbian encounter and the Little Ice Age: abrupt land use change, fire, and greenhouse forcing. *Ann. Assoc. Am. Geogr.* **100**, 755–771 (2010).
72. Nevle, R. J., Bird, D. K., Ruddiman, W. F. & Dull, R. A. Neotropical human-landscape interactions, fire, and atmospheric CO₂ during European conquest. *Holocene* **21**, 853–864 (2011).
73. Ahn, J. *et al.* Atmospheric CO₂ over the last 1000 years: a high-resolution record from the West Antarctic Ice Sheet (WAIS) divide ice core. *Glob. Biogeochem. Cycles* **26**, GB2027 (2012).
74. Rubino, M. *et al.* A revised 1000 year atmospheric delta C-13-CO₂ record from Law Dome and South Pole, Antarctica. *J. Geophys. Res.* **D 118**, 8482–8499 (2013).
75. MacFarling Meure, C. *et al.* Law Dome CO₂, CH₄ and N₂O ice core records extended to 2000 years BP. *Geophys. Res. Lett.* **33**, L14810 (2006).
76. Etheridge, D. M., Steele, L. P., Francey, R. J. & Langenfelds, R. L. Atmospheric methane between 1000 AD and present: evidence of anthropogenic emissions and climatic variability. *J. Geophys. Res.* **D 103**, 15979–15993 (1998).
77. Smith, V. C. Volcanic markers for dating the onset of the Anthropocene. *Geol. Soc. Lond. Spec. Publ.* **395**, 283–299 (2014).
78. de Silva, S. L. & Zielinski, G. A. Global influence of the AD1600 eruption of Huaynaputina, Peru. *Nature* **393**, 455–458 (1998).
79. Thompson, L. G. *et al.* Annually resolved ice core records of tropical climate variability over the past ~1800 Years. *Science* **340**, 945–950 (2013).
80. Power, M. J. *et al.* Climatic control of the biomass-burning decline in the Americas after AD 1500. *Holocene* **23**, 3–13 (2013).
81. Wang, Z., Chappellaz, J., Park, K. & Mak, J. E. Large variations in Southern Hemisphere biomass burning during the last 650 years. *Science* **330**, 1663–1666 (2010).
82. Ferretti, D. F. *et al.* Unexpected changes to the global methane budget over the past 2000 years. *Science* **309**, 1714–1717 (2005).
83. Mischler, J. A. *et al.* Carbon and hydrogen isotopic composition of methane over the last 1000 years. *Glob. Biogeochem. Cycles* **23**, GB4024 (2009).
84. Mitchell, L. E., Brook, E. J., Sowers, T., McConnell, J. R. & Taylor, K. Multidecadal variability of atmospheric methane, 1000–1800 CE. *J. Geophys. Res.* **116**, G02007 (2011).
85. Bush, M. B. & Colinvaux, P. A. Tropical forest disturbance: Paleocological records from Darien, Panama. *Ecology* **75**, 1761–1768 (1994).
86. Kinnard, C. *et al.* Reconstructed changes in Arctic sea ice over the past 1,450 years. *Nature* **479**, 509–512 (2011).
87. Neukom, R. *et al.* Inter-hemispheric temperature variability over the past millennium. *Nature Clim. Change* **4**, 362–367 (2014).
- This paper synthesizes paleoclimate records from the southern and northern hemispheres, showing one globally synchronous cool period (1594–1677) and one globally synchronous warm period (1965 onwards) within the last 1,000 years.**
88. Pomeranz, K. *The Great Divergence: China, Europe, and the Making of the Modern World Economy* (Princeton University Press, 2000).
89. Wallerstein, I. *The Modern World-System I: Capitalist Agriculture and the Origins of the European World-Economy in the Sixteenth Century* (Academic Press, 1974).
90. Killick, D. & Fenn, T. Archaeometallurgy: the study of preindustrial mining and metallurgy. *Annu. Rev. Anthropol.* **41**, 559–575 (2012).
91. Cooke, C. A., Balcom, P. H., Biester, H. & Wolfe, A. P. Over three millennia of mercury pollution in the Peruvian Andes. *Proc. Natl Acad. Sci. USA* **106**, 8830–8834 (2009).
92. Hong, S. M., Candelone, J. P., Patterson, C. C. & Boutron, C. F. History of ancient copper smelting pollution during Roman and medieval times recorded in Greenland ice. *Science* **272**, 246–249 (1996).
93. Rose, N. L. & Appleby, P. G. Regional applications of lake sediment dating by spheroidal carbonaceous particle analysis I: United Kingdom. *J. Paleolimnol.* **34**, 349–361 (2005).
94. Snowball, I., Hounslow, M. W. & Nilsson, A. Geomagnetic and mineral magnetic characterization of the Anthropocene. *Geol. Soc. Lond. Spec. Publ.* **395**, 119–141 (2014).
95. Wolfe, A. P. *et al.* Stratigraphic expressions of the Holocene-Anthropocene transition revealed in sediments from remote lakes. *Earth Sci. Rev.* **116**, 17–34 (2013).
96. Holtgrieve, G. W. *et al.* A coherent signature of Anthropogenic nitrogen deposition to remote watersheds of the Northern Hemisphere. *Science* **334**, 1545–1548 (2011).
97. Galuska, A., Migaszewski, Z. M. & Zalasiewicz, J. Assessing the Anthropocene with geochemical methods. *Geol. Soc. Lond. Spec. Publ.* **395**, 221–238 (2014).
98. Falkowski, P. *et al.* The global carbon cycle: a test of our knowledge of Earth as a system. *Science* **290**, 291–296 (2000).
99. Fairchild, I. J. & Frisia, S. Definition of the Anthropocene: a view from the underworld. *Geol. Soc. Lond. Spec. Publ.* **395**, 239–254 (2014).
100. Hua, Q. Radiocarbon: a chronological tool for the recent past. *Quat. Geochronol.* **4**, 378–390 (2009).
101. Harnisch, J. & Eisenhauer, A. Natural CF₄ and SF₆ on Earth. *Geophys. Res. Lett.* **25**, 2401–2404 (1998).
102. Butler, J. H. *et al.* A record of atmospheric halocarbons during the twentieth century from polar firn air. *Nature* **399**, 749–755 (1999).
103. Rakowski, A. Z. *et al.* Radiocarbon method in environmental monitoring of CO₂ emission. *Nucl. Instrum. Methods Phys. Res. B* **294**, 503–507 (2013).
104. Ketterer, M. E. *et al.* Resolving global versus local/regional Pu sources in the environment using sector ICP-MS. *J. Anal. At. Spectrom.* **19**, 241–245 (2004).
105. Fehn, U. *et al.* Determination of natural and anthropogenic I-129 in marine sediments. *Geophys. Res. Lett.* **13**, 137–139 (1986).
106. Hansen, V., Roos, P., Aldahan, A., Hou, X. & Possnert, G. Partition of iodine (I-129 and I-127) isotopes in soils and marine sediments. *J. Environ. Radioact.* **102**, 1096–1104 (2011).
107. Schurer, A. P., Hegerl, G. C., Mann, M. E., Tett, S. F. B. & Phipps, S. J. Separating forced from chaotic climate variability over the past millennium. *J. Clim.* **26**, 6954–6973 (2013).
108. Steffen, W., Broadgate, W., Deutsch, L., Gaffney, O. & Ludwig, C. The trajectory of the Anthropocene: the Great Acceleration. *Anthropocene Rev.* <http://dx.doi.org/10.1177/2053019614564785> (in the press).
109. Zalasiewicz, J. *et al.* When did the Anthropocene begin? A mid-twentieth century boundary level is stratigraphically optimal. *Quat. Int.* <http://dx.doi.org/10.1016/j.quaint.2014.11.045> (in the press).
110. van der Pluijm, B. Hello Anthropocene, goodbye Holocene. *Earth's Future* **2**, 2014EF000268 (2014).
111. Wright, R. A. *A Short History of Progress* (House of Anansi Press, 2004).
112. Shakun, J. D. *et al.* Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation. *Nature* **484**, 49–54 (2012).
113. Monnin, E. *et al.* Atmospheric CO₂ concentrations over the last glacial termination. *Science* **291**, 112–114 (2001).
114. Veres, D. *et al.* The Antarctic ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years. *Clim. Past* **9**, 1733–1748 (2013).
115. Marcott, S. A., Shakun, J. D., Clark, P. U. & Mix, A. C. A reconstruction of regional and global temperature for the past 11,300 years. *Science* **339**, 1198–1201 (2013).
116. Alexander, L. V. *et al.* in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker, T. F. *et al.*) 3–28 (Cambridge Univ. Press, 2013).
117. Indermuhle, A. *et al.* Holocene carbon-cycle dynamics based on CO₂ trapped in ice at Taylor Dome, Antarctica. *Nature* **398**, 121–126 (1999).
118. Siegenthaler, U. *et al.* Supporting evidence from the EPICA Dronning Maud Land ice core for atmospheric CO₂ changes during the past millennium. *Tellus B* **57**, 51–57 (2005).
119. Ahn, J. *et al.* CO₂ diffusion in polar ice: observations from naturally formed CO₂ spikes in the Siple Dome (Antarctica) ice core. *J. Glaciol.* **54**, 685–695 (2008).
120. Marín-Spiotta, E. & Sharma, S. Carbon storage in successional and plantation forest soils: a tropical analysis. *Glob. Ecol. Biogeogr.* **22**, 105–117 (2013).

121. Bonner, M. T. L., Schmidt, S. & Shoo, L. P. A meta-analytical global comparison of aboveground biomass accumulation between tropical secondary forests and monoculture plantations. *For. Ecol. Manage.* **291**, 73–86 (2013).
122. Pongratz, J., Caldeira, K., Reick, C. H. & Claussen, M. Coupled climate-carbon simulations indicate minor global effects of wars and epidemics on atmospheric CO₂ between AD 800 and 1850. *Holocene* **21**, 843–851 (2011).
123. Orihuela-Belmonte, D. E. *et al.* Carbon stocks and accumulation rates in tropical secondary forests at the scale of community, landscape and forest type. *Agric. Ecosyst. Environ.* **171**, 72–84 (2013).
124. Francey, R. J. *et al.* A 1000-year high precision record of $\delta^{13}\text{C}$ in atmospheric CO₂. *Tellus B* **51**, 170–193 (1999).
125. Trudinger, C. M., Enting, I. G., Francey, R. J., Etheridge, D. M. & Rayner, P. J. Long-term variability in the global carbon cycle inferred from a high-precision CO₂ and $\delta^{13}\text{C}$ ice-core record. *Tellus B* **51**, 233–248 (1999).
126. Böhm, F. *et al.* Evidence for preindustrial variations in the marine surface water carbonate system from coralline sponges. *Geochem. Geophys. Geosyst.* **3**, 1–13 (2002).
127. Trudinger, C. M., Enting, I. G., Rayner, P. J. & Francey, R. J. Kalman filter analysis of ice core data—2. Double deconvolution of CO₂ and $\delta^{13}\text{C}$ measurements. *J. Geophys. Res. D* **107**, D20 (2002).

Acknowledgements We acknowledge C. Hamilton, whose idea that humans are a reflexive power rather than force of nature was presented at the 'Thinking the Anthropocene' conference in Paris on 15 November 2013, and used with permission. We thank J. Kaplan and K. Krumhardt for the estimates of the population of the Americas, M. Irving for assistance with the figures, and C. Brierley, M.-E. Carr, W. Laurance, A. Mackay, O. Morton, R. Newman and C. Tzedakis for constructive discussion and remarks, and reviewer P. Gibbard for important comments. This work was funded by the European Research Council (T-FORCES, S.L.L.), a Philip Leverhulme Prize award (S.L.L.), and a Royal Society Wolfson Research Merit Award (M.A.M.).

Author Contributions S.L.L. and M.A.M. conceived the paper structure. S.L.L. conceived and developed the Obris hypothesis. S.L.L. wrote the geological importance, historical, farming and Orbis evidence reviews. M.A.M. wrote the Pleistocene, and industrialization and Great Acceleration evidence reviews. M.A.M. conceived and developed the figures. The final two sections, written by S.L.L., emerged from discussions between S.L.L. and M.A.M.

Author Information Reprints and permissions information is available at www.nature.com/reprints. The authors declare no competing financial interests. Readers are welcome to comment on the online version of the paper. Correspondence and requests for materials should be addressed to S.L.L. (s.l.lewis@ucl.ac.uk).

Copyright of Nature is the property of Nature Publishing Group and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.